

CS 450: Numerical Analysis¹

Numerical Optimization

University of Illinois at Urbana-Champaign

¹*These slides have been drafted by Edgar Solomonik as lecture templates and supplementary material for the book “Scientific Computing: An Introductory Survey” by Michael T. Heath ([slides](#)).*

Numerical Optimization

- ▶ Our focus will be on *continuous* rather than *combinatorial* optimization:

$$\min_{\mathbf{x}} f(\mathbf{x}) \quad \text{subject to} \quad \mathbf{g}(\mathbf{x}) = \mathbf{0} \quad \text{and} \quad \mathbf{h}(\mathbf{x}) \leq \mathbf{0}$$

- ▶ We consider linear, quadratic, and general nonlinear optimization problems:

Existence of Local Minima

- ▶ *Level sets* are all points for which f has a given value, *sublevel sets* are all points for which the value of f is less than a given value:

- ▶ If there exists a closed and bounded sublevel set in the domain of feasible points, then f has a global minimum in that set:

Optimality Conditions

- ▶ If \boldsymbol{x} is an interior point in the feasible domain and is a local minima,

$$\nabla f(\boldsymbol{x}) = \left[\frac{df}{dx_1}(\boldsymbol{x}) \quad \cdots \quad \frac{df}{dx_n}(\boldsymbol{x}) \right]^T = \mathbf{0} :$$

- ▶ *Critical points* \boldsymbol{x} satisfy $\nabla f(\boldsymbol{x}) = \mathbf{0}$ and can be minima, maxima, or saddle points:

Hessian Matrix

- ▶ To ascertain whether a critical point x , for which $\nabla f(x) = \mathbf{0}$, is a local minima, consider the *Hessian matrix*:

- ▶ If x^* is a minima of f , then $\mathbf{H}_f(x^*)$ is positive semi-definite:

Optimality on Feasible Region Border

- ▶ Given an equality constraint $\mathbf{g}(\mathbf{x}) = \mathbf{0}$, it is no longer necessarily the case that $\nabla f(\mathbf{x}^*) = \mathbf{0}$. Instead, it may be that directions in which the gradient decreases lead to points outside the feasible region:

$$\exists \boldsymbol{\lambda} \in \mathbb{R}^n, \quad -\nabla f(\mathbf{x}^*) = \mathbf{J}_g^T(\mathbf{x}^*)\boldsymbol{\lambda}$$

- ▶ Such *constrained minima* are critical points of the Lagrangian function $\mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}) = f(\mathbf{x}) + \boldsymbol{\lambda}^T \mathbf{g}(\mathbf{x})$, so they satisfy:

$$\nabla \mathcal{L}(\mathbf{x}^*, \boldsymbol{\lambda}) = \begin{bmatrix} \nabla f(\mathbf{x}^*) + \mathbf{J}_g^T(\mathbf{x}^*)\boldsymbol{\lambda} \\ \mathbf{g}(\mathbf{x}^*) \end{bmatrix} = \mathbf{0}$$

Sensitivity and Conditioning

- ▶ The condition number of solving a nonlinear equations is $1/f'(x^*)$, however for a minimizer x^* , we have $f'(x^*) = 0$, so conditioning of optimization is inherently bad:
- ▶ To analyze worst case error, consider how far we have to move from a root x^* to perturb the function value by ϵ :

Golden Section Search

- ▶ Given bracket $[a, b]$ with a unique minimum (f is *unimodal* on the interval), *golden section search* considers consider points $f(x_1), f(x_2)$, $a < x_1 < x_2 < b$ and discards subinterval $[a, x_1]$ or $[x_2, b]$:

- ▶ Since one point remains in the interval, golden section search selects x_1 and x_2 so one of them can be effectively reused in the next iteration:

Newton's Method for Optimization

- ▶ At each iteration, approximate function by quadratic and find minimum of quadratic function:
- ▶ The new approximate guess will be given by $x_{k+1} - x_k = -f'(x_k)/f''(x_k)$:

Safeguarded 1D Optimization

- ▶ Safeguarding can be done by bracketing via golden section search:

- ▶ Backtracking and step-size control:

Convergence of Steepest Descent

- ▶ Steepest descent converges linearly with a constant that can be arbitrarily close to 1:

- ▶ Given quadratic optimization problem $f(\mathbf{x}) = \frac{1}{2}\mathbf{x}^T \mathbf{A}\mathbf{x} + \mathbf{c}^T \mathbf{x}$ where \mathbf{A} is symmetric positive definite, the error $\mathbf{e}_k = \mathbf{x}_k - \mathbf{x}^*$ satisfies

Gradient Methods with Extrapolation

- ▶ We can improve the constant in the linear rate of convergence of steepest descent by leveraging *extrapolation methods*, which consider two previous iterates (maintain *momentum* in the direction $\mathbf{x}_k - \mathbf{x}_{k-1}$):

- ▶ The *heavy ball method*, which uses constant $\alpha_k = \alpha$ and $\beta_k = \beta$, achieves better convergence than steepest descent:

Conjugate Gradient Method

- ▶ The *conjugate gradient method* is capable of making the optimal choice of α_k and β_k at each iteration of an extrapolation method:

- ▶ *Parallel tangents* implementation of the method proceeds as follows

Krylov Optimization

- ▶ Conjugate Gradient finds the minimizer of $f(\mathbf{x}) = \frac{1}{2}\mathbf{x}^T \mathbf{A}\mathbf{x} + \mathbf{c}^T \mathbf{x}$ within the Krylov subspace of \mathbf{A} :

Newton's Method

- ▶ Newton's method in n dimensions is given by finding minima of n -dimensional quadratic approximation:

Quasi-Newton Methods

- ▶ *Quasi-Newton* methods compute approximations to the Hessian at each step:

- ▶ The *BFGS* method is a secant update method, similar to Broyden's method:

Nonlinear Least Squares

- ▶ An important special case of multidimensional optimization is *nonlinear least squares*, the problem of fitting a nonlinear function $f_{\mathbf{x}}(t)$ so that $f_{\mathbf{x}}(t_i) \approx y_i$:

- ▶ We can cast nonlinear least squares as an optimization problem and solve it by Newton's method:

Gauss-Newton Method

- ▶ The Hessian for nonlinear least squares problems has the form:
- ▶ The *Gauss-Newton* method is Newton iteration with an approximate Hessian:
- ▶ The Levenberg-Marquardt method incorporates Tykhonov regularization into the linear least squares problems within the Gauss-Newton method.

Constrained Optimization Problems

- ▶ We now return to the general case of *constrained* optimization problems:

- ▶ Generally, we will seek to reduce constrained optimization problems to a series of unconstrained optimization problems:
 - ▶ *sequential quadratic programming*:
 - ▶ *penalty-based methods*:
 - ▶ *active set methods*:

Sequential Quadratic Programming

- ▶ *Sequential quadratic programming* (SQP) corresponds to using Newton's method to solve the equality constrained optimality conditions, by finding critical points of the Lagrangian function $\mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}) = f(\mathbf{x}) + \boldsymbol{\lambda}^T \mathbf{g}(\mathbf{x})$,

- ▶ At each iteration, SQP computes $\begin{bmatrix} \mathbf{x}_{k+1} \\ \boldsymbol{\lambda}_{k+1} \end{bmatrix} = \begin{bmatrix} \mathbf{x}_k \\ \boldsymbol{\lambda}_k \end{bmatrix} + \begin{bmatrix} \mathbf{s}_k \\ \boldsymbol{\delta}_k \end{bmatrix}$ by solving

Inequality Constrained Optimality Conditions

- ▶ The *Karush-Kuhn-Tucker (KKT)* conditions hold for local minima of a problem with equality and inequality constraints, the key conditions are
- ▶ To use SQP for an inequality constrained optimization problem, consider at each iteration an *active set* of constraints:

Penalty Functions

- ▶ Alternatively, we can reduce constrained optimization problems to unconstrained ones by modifying the objective function. *Penalty* functions are effective for equality constraints $g(x) = 0$:

- ▶ The augmented Lagrangian function provides a more numerically robust approach:

Barrier Functions

- ▶ *Barrier functions* (*interior point methods*) provide an effective way of working with inequality constraints $\mathbf{h}(\mathbf{x}) \leq \mathbf{0}$: