

CS 450: Numerical Analysis¹

Partial Differential Equations

University of Illinois at Urbana-Champaign

¹*These slides have been drafted by Edgar Solomonik as lecture templates and supplementary material for the book “Scientific Computing: An Introductory Survey” by Michael T. Heath ([slides](#)).*

Partial Differential Equations

- ▶ *Partial differential equations (PDEs)* describe physical laws and other continuous phenomena:

- ▶ The *advection PDE* describes basic phenomena in fluid flow,

$$u_t = -a(t, x)u_x$$

where $u_t = \partial u / \partial t$ and $u_x = \partial u / \partial x$.

Characteristic Curves

- ▶ A *characteristic* of a PDE is a level curve in the solution:

- ▶ More generally, characteristic curves describe curves in the solution field $u(t, x)$ that correspond to solutions of ODEs, e.g. for $u_t = -a(t, x)u_x$ with $u(0, x) = u_0(x)$,

Method of Lines

- ▶ *Semidiscrete methods* obtain an approximation to the PDE by solving a system of ODEs. Consider the heat equation,

$$u_t = cu_{xx} \text{ on } 0 \leq x \leq 1, \quad u(0, x) = f(x), u(t, 0) = u(t, 1) = 0.$$

- ▶ This *method of lines* often yields a stiff ODE:

Fully Discrete Methods

- ▶ Generally, both time and space dimensions are discretized, either by applying an ODE solver to a semidiscrete method or using finite differences.
 - ▶ *Again consider the heat equation $u_t = cu_{xx}$ and discretize so $u_i^{(k)} \approx u(t_k, x_i)$,*

- ▶ *This iterative scheme corresponds to a 3-point stencil,*

Implicit Fully Discrete Methods

- ▶ Using Euler's method for the heat equation, stability requirement is
- ▶ This step-size restriction on stability can be circumvented by use of implicit time-stepper, such as backward Euler,
- ▶ Using the trapezoid method to solve the ODE we obtain the second-order *Crank-Nicolson method*,

Convergence and Stability

- ▶ *Lax Equivalence Theorem*: consistency + stability = convergence
 - ▶ *Consistency means that the local truncation error goes to zero, and is easy to verify by Taylor expansions.*
 - ▶ *Stability implies that the approximate solution at any time t must remain bounded.*
 - ▶ *Together these conditions are necessary and sufficient for convergence.*
- ▶ Stability can be ascertained by spectral or Fourier analysis:
 - ▶ *In the method of lines, we saw that the eigenvalues of the resulting ODE define the stability region.*
 - ▶ *Fourier analysis decomposes the solution into a sum of harmonic functions and bounds their amplitudes.*

CFL Condition

- ▶ The domain of dependence of a PDE for a given point (t, x) is the portion of the problem domain influencing this point through the PDE:

- ▶ *The Courant, Friedrichs, and Lewy (CFL) condition* states that for an explicit finite-differencing scheme to be stable for a hyperbolic PDE, it is necessary that the domain of the dependence of the PDE must be contained in the domain of dependence of the scheme:

Time-Independent PDEs

- ▶ We now turn our focus to time-independent PDEs as exemplified by the *Helmholtz equation*:

$$u_{xx} + u_{yy} + \lambda u = f(x, y)$$


- ▶ We discretize as before, but no longer perform time stepping:

Finite-Differencing for Poisson

- ▶ Consider the Poisson equation with equispaced mesh-points on $[0, 1]$:

Multidimensional Finite Elements

- ▶ There are many ways to define localized basis functions, for example in the 2D FEM method²:



Sparse Linear Systems

- ▶ Finite-difference and finite-element methods for time-independent PDEs give rise to sparse linear systems:
 - ▶ *typified by the 2D Laplace equation, where for both finite differences and FEM,*

- ▶ *Direct methods* apply LU or other factorization to A , while *iterative methods* refine x by minimizing $r = Ax - b$, e.g. via Krylov subspace methods.

Direct Methods for Sparse Linear Systems

- ▶ It helps to think of A as the adjacency matrix of graph $G = (V, E)$ where $V = \{1, \dots, n\}$ and $a_{ij} \neq 0$ if and only if $(i, j) \in E$:

- ▶ Factorizing the l th row/column in Gaussian elimination corresponds to removing node i , with nonzeros (new edges) introduced for each k, l such that (i, k) and (i, l) are in the graph.

Vertex Orderings for Direct Methods

- ▶ Select the node of minimum degree at each step of factorization:
- ▶ Graph partitioning also serves to bound fill, remove vertex separator $S \subset V$ so that $V \setminus S = V_1 \cup \dots \cup V_k$ become disconnected, then order V_1, \dots, V_k, S :
- ▶ *Nested dissection* ordering partitions graph into halves recursively, ordering each separator last.

Sparse Iterative Methods

- ▶ Sparse iterative methods avoid overhead of fill in sparse direct factorization.
Matrix splitting methods provide the most basic iterative methods:

Gauss-Seidel Method

- ▶ The Jacobi method takes weighted sums of $\mathbf{x}^{(k)}$ to produce each entry of $\mathbf{x}^{(k+1)}$, while Gauss-Seidel uses the latest available values, i.e. to compute $x_i^{(k+1)}$ it uses a weighted sum of

$$x_1^{(k+1)}, \dots, x_{i-1}^{(k+1)}, x_i^{(k)}, \dots, x_n^{(k)}.$$

- ▶ Gauss-Seidel provides somewhat better convergence than Jacobi:

Successive Over-Relaxation

- ▶ The *successive over-relaxation* (SOR) method seeks to improve the spectral radius achieved by Gauss-Seidel, by choosing

$$M = \frac{1}{\omega}D + L, \quad N = \left(\frac{1}{\omega} - 1\right)D - U$$

- ▶ The parameter ω in SOR controls the ‘step-size’ of the iterative method:

Conjugate Gradient

- ▶ The solution to $\mathbf{Ax} = \mathbf{b}$ when \mathbf{A} is symmetric positive definite is the minima of the quadratic optimization problem,

$$\min_{\mathbf{x}} \mathbf{x}^T \mathbf{A} \mathbf{x} - \mathbf{x}^T \mathbf{b}$$

- ▶ Conjugate gradient works by picking \mathbf{A} -orthogonal descent directions
- ▶ The convergence rate of CG is linear with coefficient $\frac{\sqrt{\kappa(\mathbf{A})}-1}{\sqrt{\kappa(\mathbf{A})}+1}$:

Preconditioning

- ▶ Preconditioning techniques choose matrix $M \approx A$ that is easy to invert and solve a modified linear system with an equivalent solution to $Ax = b$,

$$M^{-1}Ax = M^{-1}b$$

- ▶ M is chosen to be an effective approximation to A with a simple structure: